

# Markscheme

May 2023

Physics

Higher level

Paper 3

28 pages

© International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

**Subject Details: Physics HL Paper 3 Markscheme**

Candidates are required to answer **all** questions in Section A and **all** questions from **one** option in Section B. Maximum total = **45 marks**.

1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “**max**” written after the mark in the “Total” column. The related rubric, if necessary, will be outlined in the “Notes” column.
5. An alternative wording is indicated in the “Answers” column by a slash (/). Either wording can be accepted.
6. An alternative answer is indicated in the “Answers” column by “**OR**”. Either answer can be accepted.
7. An alternative markscheme is indicated in the “Answers” column under heading **ALTERNATIVE 1** etc. Either alternative can be accepted.
8. Words inside chevrons « » in the “Answers” column are not necessary to gain the mark.
9. Words that are underlined are essential for the mark.
10. The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.
11. If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
12. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
13. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. When marking, indicate this by adding **ECF** (error carried forward) on the script. “ECF acceptable” will be displayed in the “Notes” column.
14. Do **not** penalize candidates for errors in units or significant figures, **unless** it is specifically referred to in the “Notes” column.

**Section A**

Question		Answers	Notes	Total
1.	a	height «of drop» <b>OR</b> velocity «of ball» <b>OR</b> kinetic energy «of ball» <b>OR</b> temperature/mass/radius/surface area/volume of ball ✓	Allow reference to controlling spin on the ball Do <b>not</b> accept bald temperature, mass, surface area or volume.	1

Question		Answers	Notes	Total
1.	b	<p>refers to 2 non-adjacent points ✓</p> <p>suitable calculation to analyze the proportionality ✓</p> <p>identifies variation/difference in calculated values, «thus hypothesis not supported» ✓</p>	<p>Award <b>full marks</b> if more than two points used appropriately.</p> <p>Allow <b>[2 max]</b> if they use at least three points to show that two increments in force are not consistent with the corresponding increments in pressure and therefore it is not a straight line.</p>	3

Question			Answers	Notes	Total
1.	c	i	$\text{N}^2\text{m}^2$ OR $\text{kg}^2\text{m}^4\text{s}^{-4}$ OR $\text{N}^3\text{Pa}^{-1}$ ✓	<b>Award [0]</b> if they convert to base units incorrectly.	1
1.	c	ii	point plotted at (40 kPa, $49 \times 10^5 \text{ N}^3$ ) ✓	Allow for the point to be plotted from 46 to $56 \times 10^5 \text{ N}^3$ at 40 kPa, as candidates may calculate or may plot from a graphical analysis.	1
1.	d	i	15% seen anywhere ✓ « $\Delta(F^3)$ » = $39.4 \times 10^5 \times 0.15 = 5.9 \times 10^5$ ✓ $\pm 6 \times 10^5$ ✓	<b>MP1</b> is for the propagation of 5%. It can be shown differently, e.g. 3x5% Allow students to use $40 \times 10^5$ (from the graph). Award <b>MP3</b> for <b>any</b> uncertainty rounded to 1 significant digit <b>Award [3]</b> for a <b>BCA</b> . Allow <b>ECF</b> from <b>MP1</b> and <b>MP2</b>	3
1.	d	ii	error bar drawn at 30 kPa from $34 \times 10^5$ to $46 \times 10^5 \text{ N}^3$ ✓	Allow $\pm$ half square on each side of the bar or one square overall ( $\pm 2 \times 10^5$ ) Allow <b>ECF</b> from <b>d(i)</b> .	1
1.	d	iii	a «straight» line can be drawn that passes through origin ✓		1

Question		Answers	Notes	Total
2.	a	final temperature of equilibrium/water/cube <b>OR</b> mass of water/cube ✓	<p><i>Do not award mark if any additional incorrect measurement is included</i></p> <p><i>Accept temperature change if water or cube specified</i></p>	1
2.	b	smaller mass of cube <b>OR</b> hotter cube <b>OR</b> more mass of water <b>OR</b> colder water <b>OR</b> more precise thermometer ✓	<p><i>Accept reference to repeating the experiment and taking mean values.</i></p> <p><i>Accept any reasonable answer that increases change in temperature of the cube.</i></p> <p><i>Do not accept more accurate thermometer</i></p>	1
2.	c	cube specific heat will be too large/increased value/overestimate ✓ additional «thermal» energy transferred <b>OR</b> temperature rise of water will be larger <b>OR</b> temperature drop of cube will be smaller ✓		2

**Section B**

**Option A — Relativity**

Question			Answers	Notes	Total
3.	a		set of coordinates/axes to record position and time «of an event» <b>OR</b> a coordinate system which is at rest/not moving relative to the observer ✓	<i>Some mention of time <b>and</b> position for <b>MP1</b></i> <i>Allow “set of clocks and rulers” as per <b>MP1</b></i>	1
3.	b	i	magnetic ✓ «observer O sees» moving charge in a magnetic field ✓	<i>Do <b>not</b> allow electromagnetic</i> <b>OWTTE</b>	2
3.	b	ii	electric / electrostatic ✓ positive lattice ions are length contracted relative to e <b>OR</b> electrons sees positive wire ✓		2

Question		Answers	Notes	Total
4.	a	4.8 «light years» ✓		1
4.	b	«-» 0.48c ✓	Ignore sign	1
4.	c	$\gamma = 1.6$ ✓ $D = \frac{4.8}{1.6} = 3$ «ly» ✓	Allow <b>ECF</b> from <b>MP1</b>	2
4.	d	$= \frac{0.3c - 0.78c}{1 - \frac{0.78c \times 0.3c}{c^2}}$ ✓ «-» 0.63c ✓	<b>Award [2]</b> if <b>MP1</b> correct and correct answer given to 1 significant figure.	2
4.	e	$\Delta t_p = \frac{3\text{ly}}{(0.78 - 0.627)c}$ <b>OR</b> $\Delta t_p = 1.6 \left( \frac{4.8\text{ly}}{0.3c} - \frac{0.78c \times 4.8\text{ly}}{c^2} \right)$ ✓ = 19 <b>OR</b> 20 «years» ✓	<b>Award [2]</b> for <b>BCA</b>	2
4.	f	shuttle measures proper time ✓ as the events occur at the same place for the shuttle / shuttle is at both events ✓		2

Question		Answers	Notes	Total
5.	a	$0.6 c$ <b>OR</b> $= 1.8 \times 10^8 \text{ «ms}^{-1}\text{»} \checkmark$		1
5.	b	Line drawn at $45^\circ$ from $ct = 2 \text{ km}$ to hit spaceship world line at $ct = 5 \text{ km}$ <b>OR</b> $ct = 1.2/(c-0.6c) + 2 = 5 \text{ «km»} \checkmark$ $t = \frac{5000}{c} = 1.7 \times 10^{-5} \text{ «s»} \checkmark$	<i>Award [2] for BCA</i>	2
5.	c	$(ct')^2 - 0 = 5^2 - 3^2$ <b>OR</b> $\gamma = 1.25 \checkmark$ $ct' = 4 \text{ «km»}$ <b>OR</b> $t' = 13 \text{ «}\mu\text{s»} \checkmark$	<i>Allow ECF from (b)</i> <i>Allow ECF from MP1</i>	2

Question		Answers	Notes	Total
6.	a	227 «MeVc <sup>-1</sup> »		1
6.	b	$E_{\pi^+} = \sqrt{140^2 + 340^2}$ <p><b>OR</b></p> = 368 «MeV»	<i>Award [1] for correct substitution or value determined to 3 SF</i>	1
6.	c	E Pion minus = root of $(140^2 + 113^2) = \text{«180 MeV»} \checkmark$ Rest Energy of K <sup>0</sup> = $\sqrt{(368 + 180)^2 - (340 - 113)^2} \checkmark$ = 499 «MeV» ✓	<i>Award [3] for <b>BCA</b></i>	3

Question		Answers	Notes	Total
7.	a	some photon energy is used to do work against the gravitational field <b>OR</b> mention of gravitational redshift ✓	<b>OWTTE</b>	1
7.	b	$\frac{1.2 \times 10^{-4}}{3.2 \times 10^{10}} = \frac{g \times 20}{c^2}$ ✓ = 16.9 «m s <sup>2</sup> » ✓	<b>Award [2] for BCA</b>	2
7.	c	gravitational effects cannot be distinguished from inertial effects ✓ «apparent» acceleration within rocket is greater than <i>g</i> <b>OR</b> there is a stronger gravitational field <b>OR</b> the receiver is moving further away ✓	<b>OWTTE</b>	2

Option B — Engineering physics

Question		Answers	Notes	Total
8.	a	$\Sigma \Gamma = 50 \times 0.5 + 40 \times 0.2$ <b>OR</b> 33 «Nm» ✓	Accept opposite rotational sign convention	1
8.	b	$\ll \alpha = \frac{20}{5} = \gg 4 \ll \text{rad s}^{-2} \gg \checkmark$		1
8.	c	$I = \frac{\Gamma}{\alpha}$ <b>OR</b> $33 = I \times 4 \checkmark$ $I = 8.25 \ll \text{kg m}^2 \gg \checkmark$	Allow <b>ECF</b> from (a) and (b) Award <b>[2]</b> for a <b>BCA</b>	2

Question			Answers	Notes	Total
8.	d	i	moment of inertia increases ✓ Angular momentum is conserved ✓	Allow algebraic expressions e.g. $\omega = \frac{L}{I}$ so $\omega$ decreases for <b>MP2</b>	2
8.	d	ii	$E_k \llcorner = \frac{1}{2} I \omega^2 \Rightarrow \frac{1}{2} (I \omega) \omega = \frac{1}{2} L \omega \checkmark$	Accept equivalent methods	1
8.	d	iii	$\llcorner E_k \Rightarrow \frac{1}{2} L \omega_1 = \frac{1}{2} L \omega_2$ <b>OR</b> $\frac{E_{k1}}{E_{k2}} = \frac{\omega_1}{\omega_2}$ <b>OR</b> $\llcorner L$ is constant so $\llcorner E_k$ is proportional to $\omega \checkmark$ 40% $\llcorner$ energy loss $\llcorner \checkmark$	<b>MP1</b> is for understanding that angular momentum is constant so change in rotational kinetic energy is proportional to change in angular velocity <b>Award [0]</b> if $E = 0.5 I \omega^2$ is used with the same $I$ value for both values of $E$ <b>Award [2]</b> for <b>BCA</b>	2
8.	e		one example specified eg friction, air resistance, mass distribution not modelled ✓	<b>Award [1]</b> for any reasonable physical parameter that is not consistent with the model	1


Question		Answers	Notes	Total
9.	a	use of $pV = \text{constant}$ ✓ $P_B = 43 \text{ «kPa»}$ ✓	<b>Award [2] for BCA</b>	2
9.	b	concave curved line from B to locate C with a higher pressure than A ✓ vertical line joining C to A ✓	Allow <b>ECF</b> from <b>MP1</b> i.e., <b>award [1]</b> for first process locating C at a lower pressure than A, then vertical line to A.  Arrows on the processes are not needed.  Point C need not be labelled.	2
9.	c	<b>ALTERNATIVE 1</b> use of $TV^{\frac{2}{3}} = \text{constant}$ «so $300(3V_A)^{\frac{2}{3}} = T_C(V_A)^{\frac{2}{3}}$ » ✓ $T_C = 624 \text{ «K» OR } T_C = 351 \text{ «°C»}$ ✓  <b>ALTERNATIVE 2</b> use of $pV^{\frac{5}{3}}$ to get either $p_c = 43(3)^{\frac{5}{3}}$ <b>OR</b> $p_c = 268 \text{ «kPa»}$ ✓ « $T_c = 268 \times 300 / 129 = \text{so}$ » $T_C = 624 \text{ «K» OR } T_C = 351 \text{ «°C»}$ ✓		2

Question		Answers	Notes	Total
9.	d	<p><b>ALTERNATIVE 1</b> «the process is adiabatic so» <math>\Delta Q = 0</math> ✓</p> <p><b>ALTERNATIVE 2</b> The compression is reversible «so <math>\Delta S = 0</math>» ✓</p>	<b>OWTTE</b>	1
9.	e	area under curve AB is less than area under curve BC ✓	Do <b>not</b> allow <b>ECF</b> from part (b)	1
9.	f	<p>«<math>W = 0</math> so» <math>Q = \Delta U</math> ✓</p> <p>«<math>\Delta U = \frac{3}{2} \times 53.2 \times R \times (351 - 27)</math> so » <math>\Delta U = 2.15 \times 10^5</math> «J» ✓</p>	<b>Award [2] for BCA</b>	2

Question		Answers	Notes	Total
10.	a	<p>calculates <math>P_{\text{bottom}}</math> « <math>= 850 \times 9.8 \times 4 = P_{\text{bottom}}</math> « <math>= 850 \times 9.8 \times 4</math> » = 33 320 Pa »</p> <p><b>OR</b></p> <p>calculates weight of oil « <math>\text{Weight}_{\text{oil}} = 850 \times \left(\frac{0.1^2 \pi}{4}\right) \times 4 \times 9.8</math> » ✓</p> <p>Add the weight of the stopper « <math>F_{\text{min}} = 33\,320 \times \left(\frac{0.1^2 \times \pi}{4}\right) + 2.5 \times 9.8</math> » ✓</p> <p><math>F_{\text{min}} = 286 \text{ N}</math> ✓</p>	<p><i>Alternative method may include finding the volume and then weight of oil above the stopper</i></p> <p><b>Award [2 max]</b> if the weight of stopper is ignored to give 262 N</p> <p><b>Award [2 max]</b> if plug is taken as a square base to give 358 N</p> <p>Allow <b>ECF</b> from <b>MP1</b></p>	3
10.	b	<p>the pressure «at the valve opening and at the top of the oil in the tank» is constant ✓</p> <p>the velocity «of oil surface» at the top «of the tank» is zero / negligible ✓</p> <p>no energy/head losses «as the oil flows through the valve» ✓</p> <p>no turbulence/ laminar flow occurs «as the oil flows through the valve» ✓</p> <p>fluid not compressible ✓</p> <p>ideal fluid ✓</p> <p>no viscosity ✓</p>		2 max
10.	c	<p>Use of <math>\text{Re} = 1000</math> <b>OR</b> states <math>\text{Re} = \frac{\sqrt{2g(4-0.5)} \times r \times 850}{0.25}</math> ✓</p> <p>« <math>1000 = \frac{\sqrt{2g(4-0.5)} \times r \times 850}{0.25}</math> so » <math>r = 3.6</math> «cm» ✓</p>	<p><b>Award [2]</b> if <math>r</math> is taken to be the diameter this gives <math>r = 1.8 \text{ cm}</math></p>	2

Question		Answers	Notes	Total
11.	a	Q decreases so damping increases ✓ amplitude is lower «everywhere» <b>OR</b> peak shifts left / to lower frequency ✓		2
11.	b	$\frac{2\pi}{30}$ «= $\frac{\text{energy dissipated per cycle}}{\text{energy stored}}$ » = 0.21 or 21 % ✓	Accept in exact form as $\frac{2\pi}{30}$ or $\frac{\pi}{15}$	1

Option C — Imaging

Question		Answers	Notes	Total
12.	a	one ray drawn correctly reflected ✓ bottom and top of image located within the accepted region ✓	 <p>Allow the image to be within 1 square wide in the shaded region in the diagram above. The bottom not closer than 1 square from the main axis.</p>	2
12.	b	real <b>OR</b> inverted <b>OR</b> smaller image or magnification < 1 ✓	Must have any <b>two</b> correct to score [1] Features to be consistent with (a)	1
12.	c	spherical aberration <b>OR</b> rays do not meet at common focus <b>OR</b> blurred image ✓ parabolic «shape» ✓		2

Question			Answers	Notes	Total
13.	a		$M = \llcorner \frac{25}{8} \Rightarrow 3.1 \checkmark$		1
13.	b	i	$\frac{1}{8} = \frac{1}{14} + \frac{1}{v}$ <b>OR</b> $v = 18.7 \checkmark$ $M = \frac{18.7}{14}$ <b>OR</b> $M = 1.33$ <b>OR</b> $M = 1.34 \checkmark$	<i>Do not allow ECF for wrong value for v</i>	2
13.	b	ii	$\frac{1}{8} = 1 / (25-18.66) + 1/v$ <b>OR</b> $v = -30.4 \checkmark$ $m = \llcorner m_1 \times m_2 = 1.33 \times \frac{30.4}{(25 - 18.66)} = 1.33 \times 4.8 \Rightarrow 6.4 \checkmark$	<i>Allow ECF from MP1</i> <i>Allow ECF from b(i)</i>	2
13.	c	i	better resolution «than the eye» $\checkmark$	<i>Allow reference to brighter, sharper or clearer image.</i>	1
13.	c	ii	no magnification <b>OR</b> $m=1 \checkmark$		1
13.	d		international means base line can be across countries/continents $\checkmark$ greater distances increase the effective diameter of the dish $\checkmark$ great diameter improves resolution $\checkmark$	<b>OWTTE</b>	2 max

Question		Answers	Notes	Total
14.	a	attenuation = $10 \log_{10}(0.015)$ <b>OR</b> = -18.2 ✓ length = $\frac{-18.2}{-0.3}$ <b>OR</b> 60.8 «km» ✓	Allow <b>ECF</b> from <b>MP1</b> Ignore signs	2
14.	b	$\Delta t = 14 \times 10^{-5} \text{ s}$ ✓ $\left\langle \frac{27 \times 10^3}{14 \times 10^{-5}} = \frac{3 \times 10^8}{n} \right\rangle : n = 1.56$ ✓	Award <b>MP1</b> if $\Delta t = 14 \times 10^{-5} \text{ s}$ is seen anywhere Allow <b>ECF</b> for <b>MP2</b> , if candidates use $\Delta t = 18 - 3$ <b>OR</b> $\Delta t = 20 - 4$ , accepting $n = 1.67$ or $n = 1.78$	2
14.	c	refractive index of fibre is less at the edges ✓ reduces time taken for the longer path signals ✓	<b>OWTTE</b>	2

Question			Answers	Notes	Total
15.	a		$\frac{\ln 2}{0.348}$ <b>OR</b> 1.99 «cm» ✓	Allow 2.0 «cm»	1
15.	b		$\frac{I}{I_0} = e^{-0.173 \times 4} \checkmark$ $= 0.50 \checkmark$	Award [2] for BCA	2
15.	c		$I_{\text{muscle}} \times I_{\text{bone}} \times I_{\text{muscle}} = \left(\frac{1}{2}\right)^3 \checkmark$ $I = 0.125 I_0 \checkmark$	Award [2] for BCA Watch for <b>ECF</b> from (b) Allow <b>ECF</b> from <b>MP1</b>	2
15.	d		more contrast with 1 keV <b>OR</b> less sharp with 1 keV correct explanation for one of them, i.e. «more contrast with 1 keV» as coefficients have greater ratio <b>OR</b> «less sharp with 1 keV» as the X-rays will be scattered more ✓	Accept reverse argument in each case for 10 keV.	2
15.	e	i	A scan is one dimensional ✓ B scan is two dimensional ✓ B scan computed from multiple images ✓		1 max
15.	e	ii	gradient field is added to initial strong magnetic field ✓ varies linearly across/along the patient ✓ the flip/Larmor frequency varies linearly across the patient «hence position of proton frequency known» ✓		2 max

Option D — Astrophysics


Question			Answers	Notes	Total
16.	a	i	cloud/body of dust and gas ✓	<b>Award [1]</b> for a reference to dust <b>OR</b> gas. Ignore further references to origin of cloud.	1
16.	a	ii	observation of light from/passing through nebula ✓ part of the model of stellar evolution ✓	<b>MP1</b> is for any suitable comment that refers to making inferences from the radiation received, including measurement of luminosity or comparison to radiation on Earth to e.g., obtain chemical composition or wavelength shift.	1 max
16.	b	i	Star Y ✓ because parallax angle is greater <b>OR</b> star Y is closer «and that means movement relative to distant stars is greater» ✓	Allow reverse argument for star X	2
16.	b	ii	« distance = $\left(\frac{1}{0.019}\right) \times 3.26 \times 9.46 \times 10^{15}$ » $1.6 \times 10^{18}$ «m» ✓		1
16.	b	iii	$\frac{\text{Luminosity of Star X}}{\text{Luminosity of Star Y}} = \frac{b_x d_x^2}{b_y d_y^2} \checkmark$ $= 10.8 \approx 11 \checkmark$	<b>Award MP1</b> if ratio shown with distance or parallax angle. <b>Award MP1</b> for any correct substitution into ratio expression <b>Award [2]</b> for <b>BCA</b> Allow <b>ECF</b> for incorrect distances from <b>b(i)</b> or <b>b(ii)</b> .	2

Question		Answers	Notes	Total
17.	a	Hydrogen ✓		1
17.	b	stars have same/similar $L$ <b>AND</b> star B has lower $T$ ✓ correct reference to luminosity formula ( $L \propto AT^4$ ) ✓	<b>MP1</b> Allow reverse argument i.e., star A has higher $T$	2
17.	c	Any evidence of correct identification that three dots bottom left represent white dwarfs ✓ line passing through all 3 white dwarfs <b>OR</b> line continuing from 3 white dwarfs with approximately same gradient, in either direction ✓	Award <b>MP2</b> if no line drawn through the three dots but just beyond them in either direction	2
17.	d	«inward» gravitational force/pressure ✓ balanced by «neutron» degeneracy pressure/force ✓	Allow force or pressure <b>OWTTE</b>	2

Question		Answers	Notes	Total
18.	a	<p><b>ALTERNATIVE 1</b></p> <p><math>v = 0.13 \times 3 \times 10^8</math> <b>OR</b> <math>0.39 \times 10^8</math> «m s<sup>-1</sup>» <b>OR</b> <math>0.13 \times 3 \times 10^5</math> <b>OR</b> <math>0.39 \times 10^5</math> «km s<sup>-1</sup>» ✓</p> <p><math>d = \left\langle \frac{0.39 \times 10^8}{73 \times 10^3} \right\rangle \Rightarrow 530</math> «Mpc» ✓</p> <p><b>ALTERNATIVE 2</b></p> <p><math>d = \frac{cz}{H}</math> ✓</p> <p><math>\left\langle 3 \times 10^8 \times \frac{0.13}{73 \times 10^3} \right\rangle \Rightarrow 530</math> «Mpc» ✓</p>	<p><b>Award [2] for BCA</b></p>	<p><b>2</b></p>

Question			Answers	Notes	Total
18.	b	i	<p><b>ALTERNATIVE 1</b></p> $\frac{R}{R_0} = 1 + z = 1.13 \quad \checkmark$ $\frac{R}{R_0} = \frac{t}{t_0} = 1.13 \quad \checkmark$ $t_0 = \left\langle \frac{13.4 \times 10^9}{1.13} \right\rangle = 11.9 \times 10^9 \text{ «years»} \quad \checkmark$ <p><b>ALTERNATIVE 2</b></p> <p>«distance light travelled from galaxy = » <math>530 \times 10^6 \times 3.26</math> <b>OR</b> <math>1.7 \times 10^9</math> «ly» <math>\checkmark</math></p> <p>light emitted from galaxy <math>1.7 \times 10^9</math> «years ago» <math>\checkmark</math></p> <p>age when light was emitted «<math>13.4 \times 10^9 - 1.7 \times 10^9</math> =» <math>11.7 \times 10^9</math> «years ago» <math>\checkmark</math></p>	<p><b>MP1</b> can be awarded if <b>MP2</b> is clearly seen For <b>MP2</b> allow <math>H_{\text{now}} / H_{\text{then}} = 1.13</math> then <math>H_{\text{then}} = 1/t_0</math></p> <p><b>Award [3] for BCA</b></p> <p>Allow <b>ECF</b> for <b>MP3</b></p> <p>Accept <math>1.6 \times 10^{25}</math> m for <b>MP1</b></p>	3
18.	b	ii	evidence from type Ia supernovae show an accelerating universe $\checkmark$	Accept distant galaxies are further away than expected	1

Question			Answers	Notes	Total
19.	a		$M > M_j$ <b>OR</b> minimum mass for interstellar matter to collapse into a star <b>OR</b> «interstellar» potential energy > kinetic energy ✓	Do <b>not</b> penalize if they compare gravitational force with kinetic, thermal or internal energy.	1
19.	b	i	<b>ALTERNATIVE 1</b> $T \propto \frac{M}{L}$ $L \propto \frac{M}{T}$ ✓ «and L proportional to $M^{3.5}$ then» $T \propto \frac{M}{M^{3.5}}$ ✓  <b>ALTERNATIVE 2</b> L proportional $\frac{E}{T}$  <b>AND</b> L proportional $M$ ✓ «and L proportional to $M^{3.5}$ then» T proportional $\frac{M}{M^{3.5}}$ ✓	<b>MP2</b> can be awarded if <b>MP1</b> is not.	2
19.	b	ii	$\ll T_{20} M_{20}^{2.5} = T_{\text{sun}} M_{\text{sun}}^{2.5} \text{ OR } T_{20} = 10^{10} \times \left(\frac{1}{20}\right)^{2.5} \gg$ $\ll T_{20} = \gg 1.8 \times 10^9 \text{ «s» OR } 5.6 \times 10^6 \text{ «y» } \gg$ ✓	Accept 5.6 My if unit stated	1

Question		Answers	Notes	Total
20.	a	$H = \frac{73 \times 10^3}{3.26 \times 10^6 \times 9.46 \times 10^{15}} = 2.37 \times 10^{-18} \text{ «s}^{-1}\text{» } \checkmark$ $\rho_c = \frac{3 \times (2.37 \times 10^{-18})^2}{8\pi \times 6.67 \times 10^{-11}} = 1.0 \times 10^{-26} \text{ «kg m}^{-3}\text{» } \checkmark$	<p><b>Award [2] for BCA</b>                      Allow <b>ECF</b> from <b>MP1</b>  <b>Award [1] for <math>9.5 \times 10^{18}</math> or <math>9.5 \times 10^{12}</math></b></p>	2
20.	b	any irregular semicircular shape that returns to zero $\checkmark$ must pass through present and $\frac{R}{R_0} = 1 \checkmark$	<p>cosmic scale factor / R</p>  <p>Allow <b>MP2</b> for any line reasonably going through (present, 1)</p>	2
20.	c	presence of dark energy accelerates the expansion of the universe $\checkmark$ this increases the rate of cooling $\checkmark$		2