

Markscheme

May 2024

**Mathematics:
Applications and interpretation**

Higher level

Paper 2

© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- R** Marks awarded for clear **Reasoning**.
- AG** Answer given in the question and so no marks are awarded.
- FT** Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg **M1**, **A2**.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies **A3**, **M2** etc., do **not** split the marks, unless there is a note.
- The response to a “show that” question does not need to restate the **AG** line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award **FT** marks as appropriate but do not award the final **A1** in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	$8\sqrt{2}$	5.65685... (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)

2.	$\frac{35}{72}$	0.468111... (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)
----	-----------------	--	--	---

3 Implied marks

Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then **FT** marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is **(M1)A1**, it is possible to award full marks for *their* correct answer, **without working being seen**. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word “their” in a description, to indicate that candidates may be using an incorrect value.
- If the candidate’s answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any **FT** marks in the subsequent parts. This includes when candidates fail to complete a “show that” question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these **FT** rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was “Hence”.

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular misread. Use the **MR** stamp to indicate that this has been a misread and do not award the first mark, even if this is an **M** mark, but award all others as appropriate.

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- **MR** can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for parts of questions are indicated by **EITHER . . . OR**.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come “*from the use of 3 sf values*”.

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$.

An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^x$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so $x(x+1)$ and $x^2 + x$ are both acceptable.

Please note: intermediate **A** marks do NOT need to be simplified.

9 Calculators

A GDC is required for this paper, but if you see work that suggests a candidate has used any calculator not approved for IB DP examinations (eg CAS enabled devices), please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is “first”.


1. (a) 0.5

A1
[1 mark]

(b) 0.452 (0.452209...)

A2
[2 marks]

(c) (i)



A1A1

Note: Award **A1** for a normal curve (with symmetry and some evidence of change of curvature towards the extreme values).
Award **A1** for a shaded region $x < k$, where $k < \text{mean}$.

(ii) $P(T < k) = 0.3$

solving a cumulative distribution function **OR**
use of inverse function on GDC
 $k = 57.7$ (57.7071...)

(M1)
A1
[4 marks]

(d) recognizing binomial distribution
 $B(5, 0.3)$ ($P(X = 2)$)

(M1)
(A1)

0.309 (0.3087)

A1
[3 marks]

(e) $2(x-1) + 4.5$ **OR** $2x + 2.5$

A1A1

Note: Award **A1** for a linear expression with a gradient of 2,
A1 for a completely correct expression in x .

[2 marks]

(f) (\$) 13.10 (accept 13.1)

A1

[1 mark]

(g) attempt to solve $2(x-1) + 4.5 = 7.2$ **OR** $2x + 2.5 = 7.2$
2.35 (kg)

(M1)
A1

Note: Award **M1A1FT** for an answer of 1.35 (kg) from $2x + 4.5$ seen in (e).

[2 marks]

[Total 15 marks]

2. (a) (i)
$$\begin{pmatrix} 7.2 \\ 5.1 \\ 2.4 \end{pmatrix} - \begin{pmatrix} 0 \\ 0 \\ 2.8 \end{pmatrix} = \begin{pmatrix} 7.2 \\ 5.1 \\ -0.4 \end{pmatrix}$$
 A1

Note: Accept alternate vector notation, e.g. $(7.2, 5.1, -0.4)$ or $\langle 7.2, 5.1, -0.4 \rangle$

(ii) use of correct formula to find $\left| \vec{AB} \right|$ **(M1)**

$$\sqrt{7.2^2 + 5.1^2 + (-0.4)^2}$$

8.83 (km) (8.83232...)

A1
[3 marks]

(b) magnitude of $\begin{pmatrix} 1.1 \\ 8.4 \\ 0.2 \end{pmatrix}$ is

$$\sqrt{1.1^2 + 8.4^2 + 0.2^2} (= 8.47407...)$$

(A1)

EITHER

$$\begin{pmatrix} 1.1 \\ 8.4 \\ 0.2 \end{pmatrix} \cdot \begin{pmatrix} 7.2 \\ 5.1 \\ -0.4 \end{pmatrix}$$

(M1)

$$1.1 \times 7.2 + 8.4 \times 5.1 - 0.2 \times 0.4 (= 50.68)$$

(A1)

Note: The **M** mark can be implied by a partially correct **A1** line.

angle = $\arccos\left(\frac{50.68}{8.83232... \times 8.47407...}\right)$ **(M1)**

OR

Attempt to find $\begin{pmatrix} 1.1 \\ 8.4 \\ 0.2 \end{pmatrix} \times \begin{pmatrix} 7.2 \\ 5.1 \\ -0.4 \end{pmatrix}$ **(M1)**

$$\left| \begin{pmatrix} 1.1 \\ 8.4 \\ 0.2 \end{pmatrix} \times \begin{pmatrix} 7.2 \\ 5.1 \\ -0.4 \end{pmatrix} \right| = \sqrt{4.38^2 + 1.88^2 + 54.87^2} (= 55.0766...)$$
 (A1)

angle = $\arcsin\left(\frac{55.0766...}{8.83232... \times 8.47407...}\right)$ **(M1)**

THEN

47.4° (47.3805...) **OR** 0.827 (0.826947...)

A1
[5 marks]

continued...

Question 2 continued

- (c) using sum of angles in a triangle equals 180 (M1)
 $\hat{ACB} = 180 - 47.3805 - 55.2 (= 77.4194\dots)$ (A1)

$$\frac{AC}{\sin(55.2)} = \frac{8.83232\dots}{\sin(77.4194\dots)}$$

7.43 (km) (7.43107\dots) (A1)

A1
[4 marks]
[Total 12 marks]

- 3. (a) $\frac{40000}{x^2} = 400$ (M1)
 $x = 10$ (pesos) (since x is positive) A1

[2 marks]

- (b) (i) $\left(\frac{40000}{50^2} =\right) 16$ A1

- (ii) $(50 \times 16 =) 800$ (pesos) A1
[2 marks]

- (c) (i) **EITHER**
profit for each smoothie = $x - 20$ (M1)

$$P = \frac{40000}{x^2} \times (x - 20)$$

A1

- OR**
profit = revenue - costs = $nx - 20n$ (M1)

$$P = x \times \frac{40000}{x^2} - 20 \times \frac{40000}{x^2}$$

A1

Note: Do not award **A1** if $\frac{40000}{x}$ seen as first term unless explained (in part (a) or (b)), as it is given in question.

THEN

$$P = \frac{40000}{x} - \frac{800000}{x^2}$$

AG

continued...

Question 3 continued

(ii) attempt to express P ready for power rule **(M1)**

$$P = 40000x^{-1} - 800000x^{-2}$$

$$\frac{dP}{dx} = -\frac{40000}{x^2} + \frac{1600000}{x^3} \quad \text{OR} \quad \frac{dP}{dx} = -40000x^{-2} + 1600000x^{-3} \quad \text{A1A1}$$

Note: The **(M1)** can be awarded for either of the correct terms seen.
A1 for each correct term.
 At most **M1A1A0** if additional terms seen.

(iii) attempt to find x -value **(M1)**

e.g. sketch of $\frac{dP}{dx}$ with x -intercept indicated **OR** recognition that it occurs at the maximum of P **OR** algebraic approach (requires multiplication by x^3)

$$x = 40 \quad \text{A1}$$

Note: $\frac{-40000}{x^2} + \frac{1600000}{x^3} = 0$ is insufficient to award **M1**, this is given in the question. There must be an "attempt to find x -value".
 Award **M1A0** for a coordinate pair $(40, 500)$.

(iv) attempt to substitute their x -value into equation for n **(M1)**

$$n = \frac{40000}{40^2} = 25 \quad \text{A1}$$

Note: Given the nature of the function P , the local maximum is also the global maximum. This is often the case in examinations, but should not always be assumed.

[9 marks]
[Total 13 marks]

4. (a) (i) 0.9 (ii) 0.3 (iii) 0.7 **A2**

Note: Award **A1A0** if one of the values is incorrect, **A0A0** otherwise.

[2 marks]

(b) $(0.1 \times 0.3 =) 0.03$ **A1**
[1 mark]

(c) $P(\text{no fail}) = 0.63$ **(A1)**

$P(\text{one fails}) = 0.34$ **(A1)**

$P(\text{two fail}) = 0.03$ **(A1)**

Note: The three **A1**'s can be awarded independently

continued...

Question 4 continued

multiplying by 200

(M1)

No switch fails	One switch fails	Two switches fail
126	68	6

(A1)

degrees of freedom = 2

(A1)

Note: Award **A1** for df = 2 seen anywhere and may be awarded independent of the **M1** mark.
The df=2 cannot be implied from chi squared statistic = 3.40989

$$p\text{-value} = 0.182 \text{ (0.181781...)}$$

A1

$$0.182 > 0.05$$

R1

hence insufficient evidence to reject H_0 (that the manufacturers claims are correct) **A1**

Note: The **R1A1** can be awarded as follow through within part (d) from their (explicitly labelled) incorrect p-value.
An unrealistic p-value should preclude awarding the final **R1A1**.
Accept either a conclusion to not reject the null hypothesis or the manufacturers claims are correct.
Do not award **R0A1**.

[9 marks]
[Total 12 marks]

5. (a) C.

A1

Any valid reason for accepting C. or rejecting A. and B.

R1

for example:

- when $x = 0$ slopes have (or appear to have) zero gradient

- (slope field is) always positive for $x > 0$

Note: Allow **A1R0**.

[2 marks]

(b) $\int e^{2y} dy = \int x dx$

(M1)

$$\frac{1}{2}e^{2y} = \frac{1}{2}x^2 (+c)$$

(A1)(A1)

Note: **A1** for left hand side, **A1** for right hand side.

substituting in $x = 0, y = 0$

(M1)

$$\frac{1}{2} = c$$

(A1)

Note: The substitution may be seen and credited later, however at that point the constant term may be 1.

continued...

Question 5 continued

$$e^{2y} = x^2 + 1$$

$$y = \frac{1}{2} \ln(x^2 + 1)$$

M1A1

Note: Award **M1** for use of log law.

[7 marks]

(c) $\frac{dy}{dx} = \frac{1}{2} \times 2x \times \frac{1}{x^2 + 1} \left(= \frac{x}{x^2 + 1} \right)$

M1A1

Note: Award **M1** for use of chain rule, or use of implicit differentiation of the penultimate line of the answer to (b).

[2 marks]

- (d) substitution of $e^{2y} = x^2 + 1$ from part (b) into part(c)(i) or original differential equation

M1

$$\frac{dy}{dx} = \frac{x}{x^2 + 1} = \frac{x}{e^{2y}}$$

A1

and hence $y = \frac{1}{2} \ln(x^2 + 1)$ is a solution for the differential equation

AG

Note: Only award the **A1** as follow-through if their $\frac{dy}{dx}$ is of the form $\frac{x}{x^2 + c}$.

[2 marks]

[Total 13 marks]

6. (a) let S be the number of spaceships hit and B the number of battleships

- (i) mean = 8.4 **(A1)**

$P(S \leq 10) = 0.774$ (0.774301...) **A1**

- (ii) attempt to add two means **(M1)**

$4.2 + 2.3 = 6.5$

$P(S + B > 10) = P(S + B \geq 11)$ **(M1)**

0.0668 (0.0668387...) **A1**

[5 marks]

- (b) (i) $E(T) = 3 \times 4.2 + 5 \times 2.3 = 24.1$ **A1**

- (ii) $\text{Var}(T) = 3^2 \times 4.2 + 5^2 \times 2.3 = 95.3$ **(M1)A1**

[3 marks]

continued...

Question 6 continued

- (c) any valid reason **R1**
 for example:
 mean is not equal to variance **OR** T cannot take all integer values **[1 mark]**
- (d) distribution of mean score is $N\left(24.1, \frac{95.3}{60}\right)$ ($N(24.1, 1.58833\dots)$) **(A1)(A1)**

Note: Award **A1** for normal distribution with mean 24.1, and **A1** for variance $\frac{95.3}{60}$.

$P(\bar{T} > 25) = 0.238$ (0.237576...) **A2**
[4 marks]
[Total 13 marks]

7. (a) attempt to use $V = \pi \int x^2 dy$ **(M1)**
 $x^2 = 2y + 2$ or any reasonable attempt to find x in terms of y **(M1)**
 $V = \pi \int_0^h 2y + 2 dy$ **(A1)**

Note: Correct limits must be seen for the **A1** to be awarded however the dy may be omitted (as not a final answer).
 If this is given as the final answer to this part the remaining marks can be awarded if seen in part (b).

$\int 2y + 2 dy = y^2 + 2y$ **(A1)**

Note: Accept equivalent with alternate variable

$V = \pi [y^2 + 2y]_0^h$
 $= \pi(h^2 + 2h)$ **A1**

Note: The final two **A1** marks can be awarded independently of the first **A1**.
 If $h^2 + 2h$ or $y^2 + 2y$ is the final (unsupported) answer award at most **(M1)(M1)(A0)(A1)A0**.

[5 marks]

- (b) volume of vase = $\pi(15^2 + 2 \times 15)$ (= 801.106...) **(A1)**
- (time to fill vase = $\frac{801.106\dots}{20}$ =) 40.1 (40.0553...) (seconds) **A1**

Note: Accept exact answers in terms of π , e.g. 12.75π or $\frac{51\pi}{4}$

[2 marks]

continued...

Question 7 continued

(c) **EITHER**

$$\frac{dh}{dt} = \frac{dV}{dt} \times \frac{dh}{dV} \quad (M1)$$

$$\frac{dV}{dh} = \pi(2h + 2) \quad (A1)$$

OR

differentiating $V = \pi(h^2 + 2h)$ implicitly (M1)

$$\frac{dV}{dt} = \pi(2h + 2) \frac{dh}{dt} \quad (A1)$$

THEN

$$\frac{dh}{dt} = 20 \times \frac{1}{\pi(2h + 2)} \quad (M1)(A1)$$

Note: Award **M1** for attempting to solve for $\frac{dh}{dt}$, **A1** for a correct expression.

substituting $h = 10$ seen anywhere (M1)

0.289 (0.289372...) cm s⁻¹ **A1A1**

Note: Award **A1** for the correct value. Award **A1** for the correct units, independent of other marks.

[7 marks]
[Total 14 marks]

8. (a) $\begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ **M1A1**

Note: Award **M1** is for a 3x3 matrix with at least one column correct.

Column order is not explicit in question and may not be labelled in candidate response; accept their correct adjacency matrix.

[2 marks]

continued...

Question 8 continued

(b) **EITHER**

$$\begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}^5$$

(M1)

$$= \begin{pmatrix} 11 & 11 & 10 \\ 11 & 10 & 11 \\ 10 & 11 & 11 \end{pmatrix}$$

(A1)

OR

Listing at least 8 possible walks

(M1)

AAAAAB, AAABAB, AAABCB, AABAAB, AABCCB,
ABABAB, ABACAB, ABAAAB, ABCCCB, ABCBAB, ABCBCB

(A1)

THEN

11 different routes

A1
[3 marks]

(c) (i) $0.5^5 \left(\frac{1}{32}, 0.03125 \right)$

A1

(ii) **EITHER**

there are 11 possible walks so probability is 11×0.5^5

M1

OR

total number of (equally likely) walks from A is 32, 11 end up at B

M1

THEN

$\frac{11}{32}$ OR 0.344 (0.34375)

A1

Note: Solutions to this part must be using the value (11) obtained from part (b) to be awarded any marks

[3 marks]

(d) (i) $(1 \times 0.4 =) 0.4$

A1

(ii) $(0.5 \times 0.5 =) 0.25$

A1

(iii) $(0.5 \times 0.5 + 0.5 \times 0.5 =) 0.5$

A1

[3 marks]

continued...

Question 8 continued

(e) transition matrix is $\begin{pmatrix} 0 & 0.25 & 0.4 \\ 0.6 & 0.5 & 0.6 \\ 0.4 & 0.25 & 0 \end{pmatrix}$ (with order AB, AC and BC) **(M1)(A1)**

Note: Column order is not explicit in question and may not be labelled in candidate response; accept their correct transition matrix.
Accept the transposed matrix.

$$\begin{pmatrix} 0 & 0.25 & 0.4 \\ 0.6 & 0.5 & 0.6 \\ 0.4 & 0.25 & 0 \end{pmatrix}^5$$

$$= \begin{pmatrix} 0.22215 & 0.227275 & 0.23239 \\ 0.54546 & 0.54545 & 0.54546 \\ 0.23239 & 0.227275 & 0.22215 \end{pmatrix}$$

0.232 (0.23239) **(M1)**

A1
[4 marks]

- (f) (Taking a high power of a matrix)
long term probabilities are 0.227275, 0.545455 and 0.227275 **(M1)**
- B and 0.545 (54.5% $\frac{6}{11}$) **A1A1**

Note: Award **(M0)A0A0** for an unsupported answer of “B” (with either no probability or an incorrect probability).

[3 marks]
[Total 18 marks]