

© International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

# Mathematics: applications and interpretation

## Higher level

### Paper 3

9 May 2023

Zone A afternoon | Zone B morning | Zone C afternoon

1 hour

---

#### Instructions to candidates


- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Answer all the questions in the answer booklet provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics: applications and interpretation formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[55 marks]**.

Answer **both** questions in the answer booklet provided. Please start each question on a new page. Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. Solutions found from a graphic display calculator should be supported by suitable working. For example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

1. [Maximum mark: 26]

**This question considers the optimal route between two points, separated by several regions where different speeds are possible.**

Huw lives in a house, H, and he attends a school, S, where H and S are marked on the following diagram. The school is situated 1.2 km south and 4 km east of Huw’s house. There is a boundary [MN], going from west to east, 0.4 km south of his house. The land north of [MN] is a field over which Huw runs at 15 kilometres per hour ( $\text{km h}^{-1}$ ). The land south of [MN] is rough ground over which Huw walks at  $5 \text{ km h}^{-1}$ . The two regions are shown in the following diagram.



(a) Huw travels in a straight line from H to S. Calculate the time that Huw takes to complete this journey. Give your answer correct to the nearest minute. [6]

(b) Huw realizes that his journey time could be reduced by taking a less direct route. He therefore defines a point P on [MN] that is  $x$  km east of M. Huw decides to run from H to P and then walk from P to S. Let  $T(x)$  represent the time, in hours, taken by Huw to complete the journey along this route.

(i) Show that  $T(x) = \frac{\sqrt{0.4^2 + x^2} + 3\sqrt{0.8^2 + (4-x)^2}}{15}$ . [3]

(ii) Sketch the graph of  $y = T(x)$ . [2]

**(This question continues on the following page)**

**(Question 1 continued)**

(iii) Hence determine the value of  $x$  that minimizes  $T(x)$ . [1]

(iv) Find by how much Huw's journey time is reduced when he takes this optimal route, compared to travelling in a straight line from H to S. Give your answer correct to the nearest minute. [2]

(c) (i) Determine an expression for the derivative  $T'(x)$ . [3]


(ii) Hence show that  $T(x)$  is minimized when

$$\frac{x}{\sqrt{0.16 + x^2}} = \frac{3(4 - x)}{\sqrt{0.64 + (4 - x)^2}}. \quad [1]$$

(iii) For the optimal route, verify that the equation in part (c)(ii) satisfies the following result:

$$\frac{\cos \hat{H\hat{P}M}}{\cos \hat{S\hat{P}N}} = \frac{\text{speed over field}}{\text{speed over rough ground}}. \quad [2]$$

(d) The owner of the rough ground converts the southern quarter into a field over which Huw can run at  $15 \text{ km h}^{-1}$ . The following diagram shows the optimal route, HJKS, in this new situation. You are given that [HJ] is parallel to [KS].



Using a similar result to that given in part (c)(iii), at the point J, determine MJ. [6]

2. [Maximum mark: 29]

**This question considers the analysis of several datasets of examination marks using a variety of standard procedures and also an unfamiliar statistical test.**

A class of eight students sits two examinations, one in French and one in German.

The marks in these examinations are given in **Table 1**.

**Table 1**

Student	French mark	German mark
$S_1$	42	39
$S_2$	65	66
$S_3$	82	71
$S_4$	50	53
$S_5$	48	32
$S_6$	73	59
$S_7$	34	40
$S_8$	59	56

The maximum mark in both examinations is the same.

You may assume that these data are a random sample from a bivariate normal distribution with mean  $\mu_F$  for the French examination, mean  $\mu_G$  for the German examination and Pearson's product-moment correlation coefficient  $\rho$ .

Before the examinations were sat, the Head of Languages, Pierre, decided to investigate whether there would be significant evidence of a difference between  $\mu_F$  and  $\mu_G$ . He decided to analyse the data using a two-tailed paired  $t$ -test with significance level 5%.

- (a) Explain briefly
- (i) why he chose to use a  $t$ -test and not a  $z$ -test; [1]
  - (ii) why he chose to use a two-tailed test and not a one-tailed test. [1]
- (b) (i) State suitable hypotheses for the  $t$ -test. [1]
- (ii) Find the  $p$ -value for this test. [2]
  - (iii) The  $p$ -value is a probability. State the event for which it gives the probability. [1]
  - (iv) State, giving a reason, what conclusion Pierre should reach. [2]

**(This question continues on the following page)**

**(Question 2 continued)**

- (c) Pierre believes that students who score well in one language examination tend to score well in the other language examination. He therefore decides to carry out a test at the 5% significance level to investigate whether there is a positive correlation between the French examination marks and the German examination marks.
  - (i) State appropriate hypotheses in terms of  $\rho$ . [1]
  - (ii) Perform a suitable test and state the  $p$ -value. State, in context, the conclusion that Pierre should reach, giving a reason. [4]
- (d) There are actually two more students in this particular class, Paul and Sue. Paul sat the French examination, but he was unable to sit the German examination. Sue sat the German examination, but she was unable to sit the French examination.
  - (i) Paul's mark in the French examination was 58. Use the data in **Table 1** to predict the mark that Paul would have obtained in his German examination. [3]
  - (ii) Based on her mark in the German examination, Sue's mark in the French examination was predicted to be 71. Find the mark she obtained in the German examination. [2]

Six students sit examinations in mathematics and history and their marks are shown in **Table 2**. The Vice Principal, Angela, decides to investigate whether there is any association between the marks obtained in these two subjects.

**Table 2**

Student	Mathematics mark ( $x$ )	History mark ( $y$ )
$P_1$	53	41
$P_2$	76	70
$P_3$	50	62
$P_4$	65	47
$P_5$	61	66
$P_6$	84	50

**(This question continues on the following page)**

**(Question 2 continued)**

Angela is informed that the maximum mark in each subject is 100.

Angela believes that the data might not be normally distributed, so she investigates what suitable tests are available which do not assume the data is normally distributed. She decides to use an unfamiliar test based on a statistic called Kendall's  $\tau$ .

Consider  $n$  bivariate observations  $(x_i, y_i)$ ,  $i = 1, 2, \dots, n$ , such that there are no equal  $x$ -values and no equal  $y$ -values. Any pair of distinct bivariate observations  $(x_i, y_i)$  and  $(x_j, y_j)$  is said to be concordant if  $(x_i - x_j)(y_i - y_j) > 0$  and discordant if  $(x_i - x_j)(y_i - y_j) < 0$ . For  $n$  bivariate observations, there are  $\frac{n(n-1)}{2}$  distinct pairs.

Kendall's  $\tau$  is defined as  $\frac{2(C-D)}{n(n-1)}$  where  $C$  and  $D$  denote respectively the number of concordant pairs and discordant pairs.

- (e) (i) Show that the value of Kendall's  $\tau$  always lies in the interval  $[-1, +1]$ . [1]
- (ii) For students  $P_1$  and  $P_2$ , show that their pair is concordant. [1]
- (iii) Show that the value of Kendall's  $\tau$  for the mathematics and history data is 0.2. [4]
- (f) Angela decided to use this statistic in a two-tailed test at the 10% significance level. The critical region for her test is  $|\tau| \geq 0.733$ .
- (i) State, in words, her null and alternative hypotheses. [1]
- (ii) State the conclusion that Angela should reach. Give a reason for your answer. [2]

Angela now finds that the history marks are actually out of 120. The history teacher advises Angela to scale the history marks so that they are out of 100 and then redo the calculations for the value of  $\tau$ .

- (g) State, with a reason, whether you agree with the advice given by the history teacher. [2]

**References:**

© International Baccalaureate Organization 2023